

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

ASADAAME Rhinitis Drop for kids (Soda Pop)

Approved Trade Name: Asadaame kodomo bien S

You should read this written explanation before use. Please keep it so that you can read it when you need to.

Sugarless < Soda pop flavor >

Rhinitis medicine

Asadaame kodomo bien S

Class 2 OTC medicine

Precautions

When not to use the product

[If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.]

1. This drug should not be taken together with the following drugs:
Other internal medicine for rhinitis, internal medicine containing antihistamines (including cold medicine, expectorant, medicine for motion sickness, medicine for allergies), indigestion and heartburn relief.
2. After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness, blurred vision and abnormal brightness may occur).
3. Don't take this medicine for a long time.

Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.
 - (1) Patients undergoing medical treatment from a physician
 - (2) Pregnant women or women suspected of being pregnant
 - (3) The elderly
 - (4) Patients who have experienced allergic symptoms associated with drugs, etc.
 - (5) Persons with the following symptoms: Hyperthermia, Urination difficulty
 - (6) Persons diagnosed as having the following: Glaucoma, Diabetes, Thyroid disease, Heart disease, Hypertension
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Related area	Symptoms
Skin	Rash/redness, Itching
Gastrointestinal system	Nausea and vomiting, Loss of appetite
Neuropsychiatric system	Headache
Urinary system	Urination difficulty
Other	Facial hot flashes, Abnormal brightness

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

The name of the symptom	Symptoms
-------------------------	----------

Aplastic anemia	Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.
Agranulocytosis	Symptoms, such as acute high fever, chills and sore throat, etc. may occur.

- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.
Dry mouth, drowsiness, constipation, blurred vision
- When symptoms do not improve even after taking the medicine for 5 to 6 days, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

【Indications】

Relief of the following symptoms due to acute rhinitis, allergic rhinitis, or sinusitis: sneezing, runny nose, stuffy nose, watery eyes, sore throat, dull headache

【Dosage and Administration】

Take the following dose by dissolving in the mouth or by chewing.

Take this medicine in intervals of at least 4 hours.

Take this medicine in intervals of at least 7 hours.		
Age	Dose per time	Number of administrations
7 to under 15 years of age	2 tablets	3 times a day
5 to under 7 years of age	1 tablet	
Below 5 years old	Do not take	

* Do not take two tablets in your mouth at a time. Take them one at a time.

- Strictly follow the specified dosage and administration.
- When allowing children to take the medicine, a guardian must be present to watch and tell such them how to take it.

【Ingredients and Quantities】 Per 6 tablets

d-Chlorpheniramine Maleate 3mg

Phenylephrine Hydrochloride 15mg

Belladonna Total Alkaloids 0.3mg

Inactive ingredients: Acesulfame Potassium, Hydrogenated Maltose Starch Syrup, Sodium Chloride, Hydrogenated Starch Syrup, DL-Malic acid, Glycerol Esters of Fatty Acids, Acacia, Carnauba Wax, Polysorbate 80, l-Menthol, Talc, White Beeswax, Perfume.

【Precautions for storage and handling】

- Store in a cool, low humidity place with lid closed tightly, away from direct sunlight.
- Keep out of reach of children.
- Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
- Do not take this product after the expiration date. Also, take this product as soon as possible after opening the package even if the expiration date has not passed.

Manufacturer and distributor Asadaame Co., Ltd.

5-29-7, Kumegawa-Cho, Higashi-Murayama-Shi, Tokyo, Japan

Contact information: Customer service center

Phone: 03-3953-4044

Hours: 9:00 to 17:00 (excluding Saturday, Sunday, and holidays)