

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

ASADAAME Cough Drop for kids (Grape)

Approved Trade Name: Asadaame kodomo sekidome drop G

You should read this written explanation before use. Please keep it so that you can read it when you need to.

Asadaame kodomo sekidome drop G 30 tablets

Designated class 2 OTC medicine

Antitussives and Expectorants

Sugarless < Grape flavor >

Precautions

When not to use the product

[If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.]

This drug should not be taken together with the following drugs:

Other expectorants, cold medicine, and sedatives

Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.
 - (1) Patients undergoing medical treatment from a physician
 - (2) Pregnant women or women suspected of being pregnant
 - (3) Nursing women
 - (4) The elderly
 - (5) Patients who have experienced allergic symptoms associated with drugs, etc.
 - (6) Persons with the following symptoms: Hyperthermia
 - (7) Persons diagnosed as having the following: Heart disease, Hypertension, Diabetes, Thyroid disease
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Related area	Symptoms
Skin	Rash/redness, Itching
Gastrointestinal system	Nausea and vomiting, Loss of appetite
Neuropsychiatric system	Dizziness

3. When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

【Indications】

Cough, cough associated with wheezing (wheezy, whistling) , sputum, and hoarse voice due to throat inflammation, rough throat, throat discomfort, sore throat, and swollen throat.

【Dosage and Administration】

Take the following dose in the mouth and let it dissolve slowly without chewing.

Take this medicine in intervals of at least 2 hours.

Take this medicine in intervals of at least 2 hours.		
Age	Dose per time	Number of administrations
8 - 14 years	2 tablets	6 times a day
5 - 7 years	1 tablet	
Below 5 years old	Do not take	

- Strictly follow the specified dosage and administration.
- When allowing children to take the medicine, a guardian must be present to watch and tell such them how to take it.

【Ingredients and Quantities】 Per 12 tablets

dl-Methylephedrine Hydrochloride 25mg

Potassium Cresolsuphonate 90mg

Cetylpyridinium Chloride Hydrate 3mg

Inactive ingredients : Hydrogenated Starch Syrup, Hydrogenated Maltose Starch Syrup, Glycerol Esters of Fatty Acids, Dipotassium Glycyrrhizinate , Acesulfame Potassium, Monosodium L-Glutamate, DL-Malic Acid, Acacia, Carnauba Wax, Talc, White Beeswax, Perfume.

【Precautions for storage and handling】

- (1) Infants below 5 years of age should not take this product because it could be caught in the throat.
- (2) Store in a cool, low humidity place with lid closed tightly, away from direct sunlight.
- (3) Keep out of reach of children.
- (4) Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
- (5) Do not take this product after the expiration date. Also, take this product as soon as possible after opening the package even if the expiration date has not passed.

Manufacturer and distributor ASADAAME CO., LTD.

5-29-7, KUMEGAWA-CHO, HIGASHI-MURAYAMA CITY, TOKYO, JAPAN

Contact information: Customer service center

Phone: 03-3953-4044

Hours: 9:00 to 17:00 (excluding Saturday, Sunday, and holidays)